





MONTHLY CLINICAL ASSESSMENT - ACTIVE TB AND LTBI TREATMENTS

(Simultaneously refer to the tool Adverse reactions of the main TB treatments)

Starting weight: _____ kg Date: yyyy / mm / dd

EMBOSSER ICI LA CARTE DU CSI OU CSTU, SI NON DISPONIBLE, INSCRIRE LES NOM, PRÉNOM, DATE DE NAISSANCE ET NI IMÉRO DOSSIER

EMBOSS HERE THE CARD OF IHC OR UTHC, IF NOT AVAILABLE, WRITE THE NAME, SURNAME, DATE OF BIRTH AND FILE NUMBER

Presence of signs/symptoms ¹ (If abnormal, advise the physician)		y <u>yyy /nm</u> / dd		<u>yyyy /mm / dd</u>		yyyy/ mm/ dd		yyyy/ mm/ dd		<u>yyyy/mm/_dd</u>		yyyy <u>j</u> mm _/ dd	
		No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes
Hepatotoxicity ²	Changes in general condition												
	Abdominal pain/discomfort												
	Anorexia												
	Nausea/vomiting												
	Fatigue/drowsiness												
	Dark coloured urine (tea-coloured) ³												
	Pale stool (whitish)												
	Icterus/rash/pruritis												
symptoms ⁴	New or unusual cough												
	Fever												
ptc	Persistent night sweats												
	Hemoptysis												
	Weight loss (kg)												
TB	Enter the current weight ⁵		kg		kg		kg		kg		kg		kg
Other	Alcohol consumption												
	Snellen chart ⁷												
	Ishihara colour test ⁷												
	Refer to physician												
	Nurse's signature			Initials		Initials		Initials		Initials		Initials	

Initials	Signature and permit no.	Initials	Signature and permit no.	Initials	Signature and permit no.		

¹ In the presence of signs/symptoms, enter a nursing note in the record and advise the physician.

² **Hepatotoxicity:** Adults over 50 years of age, persons who drink alcohol daily, and those who have chronic liver disease are at greater risk of developing hepatotoxicity. Children can present with vague symptoms, and parents then report that they haven't been feeling well for some time.

³ Not to be confused with orange urine, which is a secondary effect of taking Rifampicin.

⁴ **TB symptoms:** New or ongoing TB symptoms can mean that the treatment is not working.

⁵ If body weight changes, have the medication adjusted.

⁶ If the answer is "Yes", enter a nursing note in the record describing the alcohol consumption and/or the test results (Snelling and Ishihara).

⁷ **If Ethambutol is taken,** perform <u>visual acuity tests</u> (Snellen chart) and <u>colour perception tests</u> (Ishihara test) q month. If results are abnormal, consult an ophthalmologist. (DSPu-TB_TB-ACT-ITL_EVAL-CLIN-MENS_EN, V2023-10-01)